

CORRECTION

Open Access



# Correction: High-fidelity elastic Green's functions for subduction zone models consistent with the global standard geodetic reference system

Takane Hori<sup>1\*†</sup> , Ryoichiro Agata<sup>1†</sup>, Tsuyoshi Ichimura<sup>2,3,4</sup>, Kohei Fujita<sup>2,4</sup>, Takuma Yamaguchi<sup>2</sup> and Takeshi Inuma<sup>1</sup>

## Correction: *Earth, Planets and Space* (2021) 73:41

<https://doi.org/10.1186/s40623-021-01370-y>

Following publication of the original article (Hori et al. 2021), the authors found that the red lines in Fig. 4, which denote the location of seismic velocity profiles in Fig. 2, is not plotted precisely. They newly plotted the lines that denote the locations of the profiles at the correct

locations in the revised version of Fig. 4. The caption of the figure was also revised accordingly. The change of the results does not affect the discussion in the main text.

The correct Fig. 4 has been provided in this Correction. The original article (Hori et al. 2021) has been corrected.

---

The original article can be found online at <https://doi.org/10.1186/s40623-021-01370-y>.

---

<sup>†</sup>Takane Hori and Ryoichiro Agata contributed equally to this study

---

\*Correspondence: horit@jamstec.go.jp


---

<sup>1</sup> Research Institute of Marine Geodynamics, Japan Agency for Marine Earth Science and Technology, Yokohama, Kanagawa, Japan

Full list of author information is available at the end of the article



© The Author(s) 2022. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.



#### Author details

<sup>1</sup>Research Institute of Marine Geodynamics, Japan Agency for Marine Earth Science and Technology, Yokohama, Kanagawa, Japan. <sup>2</sup>Earthquake Research Institute & Department of Civil Engineering, The University of Tokyo, Bunkyo-Ku, Tokyo, Japan. <sup>3</sup>Center for Advanced Intelligence Project, RIKEN, Chuo-Ku, Tokyo, Japan. <sup>4</sup>Center for Computational Science, RIKEN, Kobe, Hyogo, Japan.

Published online: 29 December 2022

#### Reference

Hori T, Agata R, Ichimura T, Fujita K, Yamaguchi T, Iinuma T (2021) High-fidelity elastic Green's functions for subduction zone models consistent with the global standard geodetic reference system. *Earth Planets Space* 73:41. <https://doi.org/10.1186/s40623-021-01370-y>

#### Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.