


CORRECTION

Open Access



Correction: Derivation of 1.064 μm normal albedos on the C-type asteroid Ryugu from laser pulse intensity measurement of the Hayabusa2 LIDAR

Ryuhei Yamada^{1*} , Keiko Yamamoto², Shoko Oshigami³, Hiroshi Araki^{2,9}, Hiroki Senshu⁴, Hiroto Noda^{5,9}, Noriyuki Namiki^{2,9}, Koji Matsumoto^{5,9}, Fumi Yoshida^{2,6}, Shinsuke Abe⁷, Naru Hirata¹, Sho Sasaki⁸ and Takahide Mizuno³

Correction: *Earth, Planets Space* (2022) 74: 166
<https://doi.org/10.1186/s40623-022-01717-z>

Following publication of the original article (Yamada et al. 2022), the authors reported an error in section “Calculation and correction of normal albedo” on page 9.

The value “0.002” was mistakenly written as “0.02” in the sentence “the frequency components from 0.02 to 0.0032 Hz are removed from the calculated albedo using low and high-pass filters” and has been corrected.

The original article (Yamada et al. 2022) has been updated.

The original article can be found online at <https://doi.org/10.1186/s40623-022-01717-z>.

*Correspondence:

Ryuhei Yamada

ryamada@u-aizu.ac.jp

¹ The University of Aizu, Ikki-Machi, Tsuruga, Aizu-Wakamatsu, Fukushima 965-8580, Japan

² National Astronomical Observatory of Japan, 2-21-1 Osawa, Mitaka, Tokyo 181-8588, Japan

³ Institute of Space and Astronautical Science, Japan Aerospace Exploration Agency, 3-1-1 Yoshinodai, Chuo-Ku, Sagami-hara, Kanagawa 252-5210, Japan

⁴ Chiba Institute of Technology, 2-17-1 Tsudanuma, Narashino, Chiba 257-0016, Japan

⁵ National Astronomical Observatory of Japan, 2-12 Hoshigaoka, Mizusawa-Ku, Oshu, Iwate 023-0861, Japan

⁶ University of Occupational and Environmental Health, 1-1 Iseigaoka, Yahatanisi-Ku, Kitakyushu, Fukuoka 807-8555, Japan

⁷ Nihon University, 7-24-1 Narashinodai, Funabashi, Chiba 274-8501, Japan

⁸ Osaka University, 1-1 Machikaneyama, Toyonaka, Osaka 560-0043, Japan

⁹ The Graduate University for Advanced Studies, SOKENDAI, Hayama, Kanagawa 240-0193, Japan

Published online: 25 January 2024

Reference

Yamada R, Yamamoto K, Oshigami S, Araki H, Senshu H, Noda H, Namiki N, Matsumoto K, Yoshida F, Abe S, Hirata N, Sasaki S, Mizuno T (2022) Derivation of 1.064 μm normal albedos on the C-type asteroid Ryugu from laser pulse intensity measurement of the Hayabusa2 LIDAR. *Earth Planets Space* 74:166. <https://doi.org/10.1186/s40623-022-01717-z>

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.